



Co všechno dávám do subarachnoidálního a epidurálního prostoru



PORODNICKÁ ANESTEZIE A ANALGEZIE – CO SE V UČEBNÍCÍCH ZATÍM NEPÍŠE



10.10.2025 11:18



ZENIT - Anestezie/perioperační medicína

Štítky: **LEKAR**

Přednášející:

Diskuzní panel expertů



OBAAMA-COV




SPINAL ANAESTHESIA (OBAAMA-COV data)

	Elective CS	Emergent CS	Elective SVK	Emergent SVK
	N=537	N=352	N=251	N=188
Position for spinal puncture				
Lying on the side	20,1 %	23,3 %	2,8 %	2,1 %
Sitting	76,4 %	72,2 %	90,4 %	91,5 %
Height of blockade application				
L1/L2	1,1 %	1,4 %	0,8 %	1,1 %
L2/L3	25,7 %	17,3 %	18,7 %	19,7 %
L3/L4	62,8 %	69,9 %	74,1 %	70,2 %
L4/L5	10,2 %	11,4 %	6,4 %	9,0 %
L5/S1	0,2 %	0,0 %	0,0 %	0,0 %
Needle				
G25	18,2 %	20,0 %	43,6 %	38,7 %
G26	48,9 %	45,1 %	27,8 %	23,7 %
G27	32,9 %	34,9 %	28,6 %	37,6 %
Introducer used	89,8 %	88,9 %	89,6 %	86,7 %
Needle tip				
Quincke/Lancet	25,6 %	33,7 %	47,2 %	53,4 %
Pencil point	48,6 %	53,0 %	38,7 %	36,2 %
Atraucan	24,2 %	12,1 %	14,0 %	10,3 %
Other	1,6 %	1,2 %	0,0 %	0,0 %
Spinal mixture				
Bupivacaine	88,3 %	84,4 %	94,8 %	91,0 %
- dose (ml) ¹	2,60 (2,40-2,80)	2,60 (2,50-2,80)	2,50 (2,20-2,60)	2,50 (2,20-2,70)
Levobupivacaine	10,8 %	15,1 %	4,4 %	10,1 %
- dose (ml) ¹	2,80 (2,50-3,00)	2,80 (2,60-3,00)	2,70 (2,50-2,95)	3,00 (2,50-3,00)
Morphine	24,4 %	24,4 %	0,0 %	0,0 %
Sufentanil	0,2 %	0,0 %	2,0 %	3,7 %
Other	0,0 %	0,0 %	0,0 %	0,0 %

Data are given in % or ¹median (IQR)

@ruthi_landau



COLUMBIA UNIVERSITY
IRVING MEDICAL CENTER

Department of
Anesthesiology


Cesarean anesthesia



Ruthi Landau

Spinal cocktail

1. Hyperbaric bupivacaine 0.75% 12mg (1.6ml)
2. Fentanyl 15 mcg (0.3ml)
3. Morphine 150 mcg (0.3ml)
4. $\pm$ clonidine 30 mcg (0.3ml)
5. $\pm$ PF dexmedetomidine 5 mcg (1.2ml)



Epidural (intrapartum)

1. Lidocaine 2% epi 20ml + HCO_3^- (or chloroprocaine 3% 20ml if emergent)
2. Fentanyl 100 mcg
3. Morphine 3 mg
4. $\pm$ clonidine 100 mcg
5. IV dexmedetomidine 10mcg

2024-12-06 12:00